

Obtaining Exact Value by Approximate Computations *

Jingzhong Zhang Yong Feng
Laboratory for Automated Reasoning and Programming
Chengdu Institute of Computer Applications
Chinese Academy of Sciences
610041 Chengdu, P. R. China
E-mail: zjz101@yahoo.com.cn, yongfeng@casit.ac.cn

Abstract

Numerical approximate computation can solve large and complex problems fast. It has the advantage of high efficiency. However it only gives approximate results, whereas we need exact results in many fields. There is a gap between approximate computation and exact results. In this paper, we build a bridge by which exact results can be obtained by numerical approximate computations.

Key words: Numerical approximate computation, Symbolic-Numerical computation, Continued fraction.

1 Introduction

Historically, the fields of symbolic computation and numerical computation have been developed by two distinct groups of people, having relatively little interaction and overlaps with each other. Symbolic computations are principally exact and stable. However they have a high complexity. Therefore, they are slow and in practice, are applicable only to small systems. Numerical approximate computation has the advantage of being fast, flexible in accuracy and being applicable to large scale problems. In Recent

*The work is partially supported by China 973 Project NKBRPC-2004CB318003. Article was submitted to Science in China in June, 2006

two decades, numerical methods are applied in the field of symbolic computations. In 1985, Kaltofen presented an algorithm for performing the absolute irreducible factorization, and suggested to perform his algorithm by floating-point numbers, then the factor obtained is an approximate one. After then, numerical methods have been studied to get approximate factors of a polynomial[4][10][14][15] [16][17]. In the meantime, numerical methods are applied to get approximate greatest common divisors of approximate polynomials [1][2][11][5], to compute functional decompositions[6], to test primality[9] and to find zeroes of a polynomial[12]. In 2000, Corless et al. applied numerical method in implicitization of parametric curves, surfaces and hypersurfaces[3]. The resulting implicit equation is still an approximate one.

There is a gap between approximate computations and exact results[18]. People usually use rational number computations to override the gap[7]. In fact, these are not approximate computations but big number computations, which are also exact computations. In 2005, Zhang et al proposed an algorithm to get exact factors of a multivariate polynomial by approximate computation[19] but they did not discuss how to override the gap. Command *convert* in maple can obtain an approximate rational number from a float if we set variable *Digits* to a positive integer. However, in order to obtain exact rational number from its approximation, we need to know two things. One is at what accuracy the float should be obtained by numerical method; another is when we should stop and return the rational number we want by the continued fraction method. So we can not obtain the exact rational number from its approximation by command *convert* and variable *Digits*. In this paper, we solve the two problems, which can be described as follows:

There is an unknown rational number m/n we want to obtain, and assume that there are approximate methods to obtain its approximation at arbitrary accuracy. We also know an upper bound N of absolute value of its denominator in advance. The two problems will be solved such as: At first, we discuss how to determine ε which is a function in N . And then use the approximate methods to obtain a floating-point number x , an approximation of m/n at accuracy $\varepsilon > 0$, i.e. $|x - m/n| < \varepsilon$. Second, we give a criteria to stop our program and return the exact rational number we want by continued fraction method.

The remainder of the paper is organized as follows. Section 2 gives a

review of continued fraction. Section 3 discusses how small the error needs to ensure the exact number to be obtained and how to get the exact number from its approximation. Section 4 gives some experimental results. The final section makes conclusions.

2 Continued fraction

A continued fraction representation of a real number is one of the forms:

$$a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{a_3 + \dots}}}, \quad (1)$$

where a_0 is an integer and $a_1, a_2, a_3, \dots$ are positive integers. One can abbreviate the above continued fraction as $[a_0; a_1, a_2, \dots]$. For finite continued fractions, note that

$$[a_0; a_1, a_2, \dots, a_n, 1] = [a_0; a_1, a_2, \dots, a_n + 1].$$

So, for every finite continued fraction, there is another finite continued fraction that represents the same number. Every finite continued fraction is rational number and every rational number can be represented in precisely two different ways as a finite continued fraction. The other representation is one element shorter, and the final term must be greater than 1 unless there is only one element. However, every infinite continued fraction is irrational, and every irrational number can be represented in precisely one way as an infinite continued fraction. An infinite continued fraction representation for an irrational number is mainly useful because its initial segments provide excellent rational approximations to the number. These rational numbers are called the convergents of the continued fraction. Even-numbered convergents are smaller than the original number, while odd-numbered ones are bigger. If successive convergents are found, with numerators $h_1, h_2, \dots$, and denominators $k_1, k_2, \dots$, then the relevant recursive relation is:

$$h_n = a_n h_{n-1} + h_{n-2}, \quad k_n = a_n k_{n-1} + k_{n-2}.$$

The successive convergents are given by the formula

$$\frac{h_n}{k_n} = \frac{a_n h_{n-1} + h_{n-2}}{a_n k_{n-1} + k_{n-2}},$$

where $h_{-1} = 1$, $h_{-2} = 0$, $k_{-1} = 0$ and $k_{-2} = 1$. Here are some useful theorems[8]:

Theorem 1 *For any positive $x \in R$, it holds that*

$$[a_0, a_1, \dots, a_{n-1}, x] = \frac{xh_{n-1} + h_{n-2}}{xk_{n-1} + k_{n-1}} \quad (2)$$

Theorem 2 *The convergents of $[a_0, a_1, a_2, \dots]$ are given by*

$$[a_0, a_1, \dots, a_n] = \frac{h_n}{k_n}$$

and

$$k_n h_{n-1} - k_{n-1} h_n = (-1)^n.$$

Theorem 3

$$\left| \frac{h_n}{k_n} - \frac{h_{n-1}}{k_{n-1}} \right| = \frac{1}{k_n k_{n-1}}$$

and

$$\frac{1}{k_n(k_{n+1} + k_n)} < \left| x - \frac{h_n}{k_n} \right| < \frac{1}{k_n k_{n+1}}$$

In order to recover exact rational number, we introduce a controlling error into the conventional continued fraction method. The continued fraction method is modified as follows.

Algorithm 1 *Continued fraction method*

Input: a nonnegative floating-point number a and $\varepsilon > 0$;

Output: a rational number b .

Step 1: $i := 1$ and $x_1 := a$;

Step 2: Getting integral part of x_i and assigning it to a_i , assigning its remains to b_i . If $b_i < \varepsilon$, then goto Step 5;

Step 3: $i := i + 1$;

Step 4: $x_i := \frac{1}{b_{i-1}}$ and goto Step 2;

Step 5: Computing expression (1) and assigning it to b .

Step 6: return b .

We will discuss the controlling error ε in algorithm 1 in the next section.

3 Recovering the exact number from its approximation

In this section, we will solve such a problem: for a given floating number w which is an approximation of rational number $\frac{m}{n}$, how do we obtain integer m and n ? Without loss of generality, we always assume that m, n are positive number. At first, we have a lemma as follows:

Lemma 1 *m, n, p, q are integer, and $pn > 0$. If $|\frac{m}{n} - \frac{q}{p}| < \frac{1}{pn}$, then $\frac{m}{n} = \frac{q}{p}$.*

Proof. $|\frac{m}{n} - \frac{q}{p}| = \frac{|pm - qn|}{pn}$. Noticing $|pm - qn|$ is a nonnegative integer, and $|\frac{m}{n} - \frac{q}{p}| < \frac{1}{pn}$, yields $|pm - qn| < 1$. Hence $|pm - qn| = 0$. That is $\frac{m}{n} = \frac{q}{p}$. The proof is finished.

Corollary 1 *m, n, p, q are integers and $p > 0, n > 0$. Let $N \geq \max\{p, n, 2\}$. If $|\frac{m}{n} - \frac{q}{p}| < \frac{1}{N(N-1)}$, then $\frac{m}{n} = \frac{q}{p}$.*

Proof: When $p \neq n$, it holds that $pn \leq N(N-1)$. Hence $|\frac{m}{n} - \frac{q}{p}| < \frac{1}{N(N-1)} \leq \frac{1}{pn}$. According to lemma 1, it is obtained that $\frac{m}{n} = \frac{q}{p}$. When $p = n$, we have

$$|\frac{m}{n} - \frac{q}{p}| = \frac{|m - q|}{n} < \frac{1}{N(N-1)} \Rightarrow |m - q| < \frac{n}{N(N-1)} \leq \frac{1}{N-1} \leq 1$$

So, $m = q$. The proof of the corollary is finished.

And now, we study how small the error $|w - \frac{m}{n}|$ needs so as to get exact rational number $\frac{m}{n}$ from its approximation w . The following theorem answers this question.

Theorem 4 *Let $x = \frac{m}{n}$ be a reduced proper fraction, and $N \geq \max\{n, 2\}$. Assume that $|x - w| < 1/(2N(N-1))$. If we get positive rational number p/q such that $|p/q - w| < 1/(2N(N-1))$, where $q \leq N$, then it holds that $x = q/p$.*

Proof: From the assumption of the theorem, we have $|x - q/p| < 1/(N(N-1))$. According to corollary 1, it holds that $q/p = m/n = x$. The proof of the theorem is finished.

We have answered how small the error is so that we can recover the exact rational number from its approximation. The remaining problem is how to get the exact number. We attack it by continued fraction method.

Let n_2/n_1 be a rational number and r_0 its approximation. Their continued fraction representations are $n_2/n_1 = [a_0, a_1, \dots, a_L]$ and $r_0 = [b_0, b_1, \dots, b_M]$ respectively. We wish that $a_i = b_i$ for $i = 0, 1, 2, \dots, L-1$ and for the last term of the continued fraction representations of n_2/n_1 , either $a_L = b_L$ or $a_L - 1 = b_L$, so that we can get n_2/n_1 from $[b_0, b_1, \dots, b_{L+1}]$. This is the following theorem:

Theorem 5 *Let n_2/n_1 be a rational number and r_0 its approximation. Assume that n_2, n_1 are coprime positive numbers, where $n_2 < n_1$, and $n_1 > 1$. The representations of n_2/n_1 and r_0 are $[a_0, a_1, \dots, a_L]$ and $[b_0, b_1, \dots, b_M]$ respectively. If $|r_0 - n_2/n_1| < 1/(4n_1(n_1 - 1))$, then one of the following statements must hold.*

- $a_i = b_i$ ($i = 0, 1, \dots, L$);
- $a_i = b_i$ ($i = 0, 1, \dots, L-1$), $a_L - 1 = b_L$, and $b_{L+1} = 1$.

According to assumption of $n_2 < n_1$, we have that $a_0 = 0$, and $b_0 = 0$. Hence $a_0 = b_0$. In order to finish the proof of theorem 5, we need two lemmas. Due to $n_2/n_1 = [a_0, a_1, \dots, a_L]$ and $r_0 = [b_0, b_1, \dots, b_M]$, we have the following expansions:

$$\frac{n_1}{n_2} = a_1 + \frac{n_3}{n_2}, \frac{n_2}{n_3} = a_2 + \frac{n_4}{n_3}, \dots, \frac{n_{L-1}}{n_L} = a_{L-1} + \frac{1}{n_L}, n_L = a_L \quad (3)$$

and

$$\frac{1}{r_0} = b_1 + r_1, \frac{1}{r_1} = b_2 + r_2, \dots, \frac{1}{r_{L-1}} = b_L + r_L, \dots, \frac{1}{r_{M-1}} = b_M \quad (4)$$

Denoting $d_i = r_i - n_{i+2}/n_{i+1}$, we have a lemma as follows:

Lemma 2 *Let n_2/n_1 be a rational number and r_0 its approximation. Assume that n_2, n_1 are coprime positive integers, where $n_2 < n_1$, and $n_1 > 1$. The representations of n_2/n_1 and r_0 are $[a_0, a_1, \dots, a_L]$ and $[b_0, b_1, \dots, b_M]$ respectively. And assume that $a_i = b_i$ for $i \leq k < L$ (k is a positive integer). Then when $|d_k| < \frac{1}{n_{k+1}(n_{k+1}-1)}$, it holds that $a_{k+1} = b_{k+1}$ for $k < L-1$; when $|d_{L-1}| < \frac{1}{n_L(n_L+1)}$, it holds that $a_L = b_L$ or $a_L - 1 = b_L$.*

Proof: At first, we show that under the assumption of the lemma if we have

$$\left| \frac{n_{k+1}^2 d_k}{n_{k+2}(n_{k+2} + n_{k+1} d_k)} \right| < \frac{1}{n_{k+2}} \quad (5)$$

then, it holds that $a_{k+1} = b_{k+1}$ for $k < L - 1$, and $a_{k+1} = b_{k+1}$ or $a_{k+1} - 1 = b_{k+1}$ for $k = L - 1$. We discuss it in two cases:

Case 1($k < L - 1$): From $d_k = r_k - n_{k+2}/n_{k+1}$, it holds that $r_k = n_{k+2}/n_{k+1} + d_k$. Hence we have that

$$\begin{aligned} \frac{1}{r_k} - \frac{n_{k+1}}{n_{k+2}} &= -\frac{n_{k+1}^2 d_k}{n_{k+2}(n_{k+2} + n_{k+1} d_k)} \Rightarrow \frac{1}{r_k} = \frac{n_{k+1}}{n_{k+2}} - \frac{n_{k+1}^2 d_k}{n_{k+2}(n_{k+2} + n_{k+1} d_k)} \\ \Rightarrow \frac{1}{r_k} &= a_{k+1} + \frac{n_{k+3}}{n_{k+2}} - \frac{n_{k+1}^2 d_k}{n_{k+2}(n_{k+2} + n_{k+1} d_k)} = b_{k+1} + r_{k+1} \end{aligned}$$

Hence, it is obvious that $a_{k+1} = b_{k+1}$ if and only if

$$0 \leq \frac{n_{k+3}}{n_{k+2}} - \frac{n_{k+1}^2 d_k}{n_{k+2}(n_{k+2} + n_{k+1} d_k)} < 1. \quad (6)$$

Therefore, if inequality (5) holds, then above inequality is guaranteed.

Case 2:(when $k = L - 1$) We have

$$\frac{1}{r_{L-1}} = a_L - \frac{n_L^2 d_{L-1}}{n_{L+1}(n_{L+1} + n_L d_{L-1})} = b_L + r_L$$

From the above equation, if

$$\left| \frac{n_L^2 d_{L-1}}{n_{L+1}(n_{L+1} + n_L d_{L-1})} \right| < \frac{1}{n_{L+2}} = 1$$

, then $a_L = b_L$ for $d_{L-1} < 0$, and $a_L - 1 = b_L$ for $d_{L-1} \geq 0$. Therefore, we have shown that if inequality (5) holds, then $a_{k+1} = b_{k+1}$ for $k < L - 1$, and $a_{k+1} = b_{k+1}$ or $a_{k+1} - 1 = b_{k+1}$ for $k = L - 1$.

On the other hand, we have

$$\begin{aligned} \left| \frac{n_{k+1}^2 d_k}{n_{k+2}(n_{k+2} + n_{k+1} d_k)} \right| &= \frac{n_{k+1}^2 |d_k|}{|n_{k+2}(n_{k+2} + n_{k+1} d_k)|} \\ &= \frac{n_{k+1} |d_k|}{n_{k+2} \left(\left| \frac{n_{k+2}}{n_{k+1}} + d_k \right| \right)} \leq \frac{n_{k+1} |d_k|}{n_{k+2} \left(\left| \frac{n_{k+2}}{n_{k+1}} - |d_k| \right| \right)} \end{aligned}$$

So, in order to ensure inequality (5), we only need it holds that

$$\frac{n_{k+1}|d_k|}{n_{k+2}|\left(\frac{n_{k+2}}{n_{k+1}} - |d_k|\right)|} < \frac{1}{n_{k+2}} \quad (7)$$

Solving inequality (7) yields

$$|d_k| < \frac{n_{k+2}}{n_{k+1}(n_{k+1} + 1)} \quad (8)$$

When $k < L - 1$, we have that $n_{k+2} > 1$. So, it holds that

$$\frac{1}{n_{k+1}(n_{k+1} - 1)} \leq \frac{n_{k+2}}{n_{k+1}(n_{k+1} + 1)}$$

Accordingly, it is obtained that

$$|d_k| < \frac{1}{n_{k+1}(n_{k+1} - 1)} \quad (9)$$

When $k = L - 1$, we have that $n_{L+1} = 1$, so it is obtained that

$$|d_{L-1}| < \frac{1}{n_L(n_L + 1)} \quad (10)$$

The proof of lemma 2 is finished.

Lemma 3 *Let n_2/n_1 be a rational number and r_0 its approximation, where n_2, n_1 are coprime positive integers, and $n_2 < n_1$, and $n_1 > 1$. The continued fraction representations of n_2/n_1 and x are $[a_0, a_1, \dots, a_L]$ and $[b_0, b_1, \dots, b_M]$ respectively. Denote $d_i = r_i - n_{i+2}/n_{i+1}$ for $i = 0, \dots, L$. Assume that $a_i = b_i$ for $i \leq k < L - 1$ (k is a positive integer). Then when $|d_k| < \frac{1}{n_{k+1}(n_{k+1}-1)}$, it holds that*

$$|d_{k+1}| < \frac{n_{k+1}(n_{k+1} - 1)}{n_{k+2}(n_{k+2} - 1)} |d_k| \quad (11)$$

Proof: Under the assumption that $a_i = b_i$ for $i = 0, 1, \dots, k$, from equation (5), we get $d_{k+1} = -\frac{n_{k+1}^2 d_k}{n_{k+2}(n_{k+2} + n_{k+1} d_k)}$. Hence we deduce a relation as follows:

$$|d_{k+1}| = \frac{n_{k+1}^2 |d_k|}{n_{k+2}^2 + n_{k+1} n_{k+2} d_k} = \frac{n_{k+1} |d_k|}{n_{k+2} \left(\frac{n_{k+2}}{n_{k+1}} + d_k \right)}$$

$$\begin{aligned}
&= \frac{n_{k+1}(n_{k+1} - 1)|d_k|}{n_{k+2}\left(\frac{n_{k+2}(n_{k+1} - 1)}{n_{k+1}} + (n_{k+1} - 1)d_k\right)} \\
&= \frac{n_{k+1}(n_{k+1} - 1)|d_k|}{n_{k+2}(n_{k+2} - 1 + \frac{n_{k+1} - n_{k+2}}{n_{k+1}} + (n_{k+1} - 1)d_k)}
\end{aligned}$$

When $|d_k| < \frac{1}{n_{k+1}(n_{k+1} - 1)}$, it holds that $\frac{n_{k+1} - n_{k+2}}{n_{k+1}} + (n_{k+1} - 1)d_k > 0$. Hence we have a relation between d_{k+1} and d_k :

$$|d_{k+1}| < \frac{n_{k+1}(n_{k+1} - 1)}{n_{k+2}(n_{k+2} - 1)}|d_k|$$

The proof of the lemma is finished.

And now, let us prove the theorem. If $|d_0| = |r_0 - n_2/n_1| < 1/(4n_1(n_1 - 1))$, From lemma 3, we can get $|d_i| < 1/(4n_{i+1}(n_{i+1} - 1))$ for $i = 0, \dots, L - 1$. Note that $n_L > n_{L+1} = 1$ and

$$\frac{1}{4n_{i+1}(n_{i+1} - 1)} < \frac{1}{n_{i+1}(n_{i+1} + 1)} < \frac{1}{n_{i+1}(n_{i+1} - 1)}$$

when $n_{i+1} > 1$. So, it holds that

$$|d_i| < \frac{1}{4n_{i+1}(n_{i+1} - 1)} < \frac{1}{n_{i+1}(n_{i+1} + 1)}$$

for $i = 0, \dots, L - 1$. According to lemma 2, the proof of the theorem is finished.

For an unknown rational number n_2/n_1 and its approximation r_0 , theorem 5 shows that $n_2/n_1 = [b_0, \dots, b_L]$ or $n_2/n_1 = [b_0, b_1, \dots, b_L, 1]$ when $|r_0 - n_2/n_1| < 1/(4n_1(n_1 - 1))$. However, we do not know what the number L is. If we make b_{L+1} large enough when $n_2/n_1 = [b_0, \dots, b_L]$, or make b_{L+2} large enough when $n_2/n_1 = [b_0, b_1, \dots, b_L, 1]$, then we recover n_2/n_1 easily. The following theorem solve this problem.

Theorem 6 *Let n_2/n_1 be a rational number and r_0 its approximation. Assume that n_2, n_1 are coprime positive integers, where $n_2 < n_1$, and $n_1 > 1$. K is a positive integer. The continued fraction representations of n_2/n_1 and r_0 are $[a_0, a_1, \dots, a_L]$ and $[b_0, b_1, \dots, b_M]$ respectively. If $|d_0| = |r_0 - n_2/n_1| < 1/((2K + 2)n_1(n_1 - 1))$, then one of the following two statements must hold*

- $a_i = b_i$ for $i = 0, \dots, L$, and $b_{L+1} \geq K$;

- $a_i = b_i$ for $i = 0, \dots, L-1$, and $b_L = a_L - 1$, $b_{L+1} = 1$, $b_{L+2} \geq K$.

Proof: From equation (3) and equation (4), we have that $\frac{n_{L-1}}{n_L} = a_{L-1} + \frac{1}{n_L}$, $n_L = a_L$ and $\frac{1}{r_{L-1}} = b_{L-1} + r_{L-1}$. When $|d_0| = |r_0 - n_2/n_1| < 1/((2K+2)n_1(n_1-1)) < 1/(4n_1(n_1-1))$, from lemma 3, it holds that $|d_i| < 1/((2K+2)n_{i+1}(n_{i+1}-1))$ for $i = 0, 1, \dots, L-1$. Furthermore, from theorem 5 it holds that $a_{L-1} = b_{L-1}$ and $d_{L-1} = r_{L-1} - \frac{1}{n_L}$. So we have

$$\frac{1}{r_{L-1}} = a_L - \frac{n_L^2 d_{L-1}}{1 + n_L d_{L-1}}$$

We discuss it in two cases:

Case 1 ($d_{L-1} > 0$): Due to $-\frac{n_L^2 d_{L-1}}{1 + n_L d_{L-1}} < 0$, we have

$$\frac{1}{r_{L-1}} = a_L - 1 + (1 - \frac{n_L^2 d_{L-1}}{1 + n_L d_{L-1}}) = b_L + r_L$$

Hence

$$\begin{aligned} r_L &= (1 - \frac{n_L^2 d_{L-1}}{1 + n_L d_{L-1}}) = \frac{1 + n_L d_{L-1} - n_L^2 d_{L-1}}{1 + n_L d_{L-1}} \\ \Rightarrow \frac{1}{r_L} &= \frac{1 + n_L d_{L-1}}{1 + n_L d_{L-1} - n_L^2 d_{L-1}} \end{aligned}$$

It is obvious that $1 < 1/r_L < 2$ when $|d_{L-1}| < 1/(4n_L(n_L-1))$. Therefore, we have

$$\frac{1}{r_L} = 1 + \frac{1 + n_L d_{L-1}}{1 + n_L d_{L-1} - n_L^2 d_{L-1}} - 1 = 1 + \frac{n_L^2 d_{L-1}}{1 + n_L d_{L-1} - n_L^2 d_{L-1}}$$

We get that $b_{L+1} = 1$ and

$$\frac{1 + n_L d_{L-1} - n_L^2 d_{L-1}}{n_L^2 d_{L-1}} = b_{L+2} + r_{L+2}. \quad (12)$$

So,

$$b_{L+2} = \left[\frac{1 + n_L d_{L-1} - n_L^2 d_{L-1}}{n_L^2 d_{L-1}} \right]$$

where $[.]$ stands for getting the integral part of a number.

We hope that b_{L+2} is greater than some integer K , which is used as a sign that $b_0, \dots, b_L, 1$ have been obtained. Solve the following inequality:

$$\begin{aligned} \frac{1 + n_L d_{L-1} - n_L^2 d_{L-1}}{n_L^2 d_{L-1}} &> K \Leftrightarrow 1 + n_L d_{L-1} - n_L^2 d_{L-1} > K n_L^2 d_{L-1} \\ \Leftrightarrow d_{L-1} &< \frac{1}{(K+1)n_L^2 - n_L} \end{aligned}$$

Since $(2K+2)n_L(n_L-1) > (K+1)n_L^2 - n_L$, we take

$$d_{L-1} < \frac{1}{(2K+2)n_L(n_L-1)}$$

Case 2($d_{L-1} < 0$): Due to $-\frac{n_L^2 d_{L-1}}{1+n_L d_{L-1}} > 0$, we have

$$\frac{1}{r_{L-1}} = a_L + \left(-\frac{n_L^2 d_{L-1}}{1+n_L d_{L-1}}\right) = b_L + r_L$$

So, $a_L = b_L$ and

$$r_L = \left(-\frac{n_L^2 d_{L-1}}{1+n_L d_{L-1}}\right)$$

It is obtained that

$$\frac{1}{r_L} = -\frac{1+n_L d_{L-1}}{n_L^2 d_{L-1}} = b_{L+1} + r_{L+1}$$

Therefore

$$b_{L+1} = \left[-\frac{1+n_L d_{L-1}}{n_L^2 d_{L-1}}\right]$$

Solve the following inequality

$$\begin{aligned} -\frac{1+n_L d_{L-1}}{n_L^2 d_{L-1}} &> K \Leftrightarrow 1+n_L d_{L-1} > n_L^2 (-d_{L-1})K \\ \Leftrightarrow (-d_{L-1}) &< \frac{1}{K n_L^2 + n_L} \end{aligned}$$

It is obvious that $(2K+2)n_L(n_L-1) > K n_L^2 + n_L$ for $n_L \geq 2$. So we take

$$-d_{L-1} < \frac{1}{(2K+2)n_L(n_L-1)}$$

The proof of theorem 6 is finished.

For practical purpose, we hope the restriction on $n_1 > 1$ and $n_1 > n_2$ can be lifted. So we have following theorem:

Theorem 7 Let n_0/n_1 be a reduced rational number and r its approximation. Assume that n_0, n_1 are positive integers and $N \geq \max\{n_1, 2\}$. K is a positive integer. The continued fraction representations of n_0/n_1 and r are $[a_0, a_1, \dots, a_L]$ and $[b_0, b_1, \dots, b_M]$ respectively. If $|d| = |r - n_0/n_1| < 1/((2K+2)N(N-1))$, then one of the following two statements must hold

- $a_i = b_i$ for $i = 0, \dots, L$, and $b_{L+1} \geq K$;
- $a_i = b_i$ for $i = 0, \dots, L-1$, and $b_L = a_L - 1$, $b_{L+1} = 1$, $b_{L+2} \geq K$.

Proof: We prove the theorem in three cases:

Case 1 ($n_1 > 1$, $n_0 < n_1$): This is theorem 6.

Case 2 ($n_1 = 1$): We have that $a_0 = n_0/n_1$. If $d = r - n_0/n_1 > 0$, then $b_0 = a_0$ and $r_0 < 1/((2K+2)2(2-1))$. So, it holds that $1/r_0 > (2(2K+2))$. Therefore, it is obtained that $b_1 > K$. If $d = r - n_0/n_1 < 0$, then

$$\begin{aligned}
r &= a_0 - |d| = a_0 - 1 + 1 - |d| = b_0 + 1 - |d| \Rightarrow r_0 = 1 - |d| \\
&\Rightarrow 1/r_0 = 1 + \frac{|d|}{1 - |d|} = b_1 + \frac{|d|}{1 - |d|} \\
&\Rightarrow 1/r_1 = \frac{1 - |d|}{|d|} = \frac{1}{|d|} - 1 > (4K + 4) - 1 \\
&\Rightarrow b_2 > 1/r_1 - 1 > 4K + 2 > K
\end{aligned}$$

So, we have that $b_0 = a_0 - 1$, $b_1 = 1$ and $b_2 > K$.

Case 3 ($n_0 > n_1$): From $n_0/n_1 = a_0 + n_2/n_1$, it holds that $n_0/n_1 - a_0 = n_2/n_1$. On the other hand, we have that $|n_0/n_1 - r| < 1/((2K+2)N(N-1)) \leq 1/n_1$. So, we can deduce that $a_0 < r < a_0 + 1$. Accordingly, it holds that $b_0 = a_0$. Hence, we have

$$\begin{aligned}
|d| &= |r - n_0/n_1| = |b_0 + r_0 - a_0 - n_2/n_1| = |r_0 - n_2/n_1| \\
&= d_0 < 1/((2K+2)N(N-1)) < 1/((2K+2)n_1(n_1-1))
\end{aligned}$$

Since $n_1 > 1$ and $n_2 < n_1$, from theorem 6, the theorem holds. Therefore, the proof is finished.

From theorem 7, we can get exact non-negative number n_2/n_1 from its approximation r as follows:

Step 1: estimating an upper bound of the denominator of n_2/n_1 , Denoted by N ;

Step 2: computing

$$d = \frac{1}{(2N + 2)N(N - 1)}$$

Step 3: obtaining r by approximate method such that $|r - n_2/n_1| < d$;

Step 4: taking $\varepsilon = 1/N$ in algorithm 1 and calling algorithm 1 to get
 b. So $b = n_2/n_1$.

4 Experimental results

The following examples run in the platform of Maple 10 and PIV 3.0G, 512M RAM. They take little time for obtaining exact rational numbers from their approximations, so we do not show time.

Example 1. Let a be unknown rational number. We only know a bound of its denominator $N = 170$. According to theorem 7, Computing rational number a as follows: Take $K = 170 + 1$, $d = (2 * K + 2) * N * (N - 1) = 9883120$, and compute $1/d = 1/9883120$. Assume that we use some numerical method to get an approximation $b = 0.8106507864$ such that $|a - b| < 1/d$. Taking $\varepsilon = 1/K$, we recover number a by algorithm 1. We get $[0, 1, 4, 3, 1, 1, 4,]$ by the first 7 steps. When doing at step 8, we get 314, which is larger than K . We stop and return $a = [0, 1, 4, 3, 1, 1, 4]$. It is $137/169$.

Example 2. Let a be unknown rational number. We only know a bound of its denominator $N = 1790$. According to theorem 7, Computing rational number a as follows: Take $K = 1790 + 1$, yields $d = (2 * K + 2) * N * (N - 1) = 11477079040$. Assume that we use some numerical method to get an approximation $b = 0.1788708777$ such that $|a - b| < 1/d$. Taking $\varepsilon = 1/K$, we recover number a by algorithm 1. We get $[0, 5, 1, 1, 2, 3, 1, 6, 2,]$ by the first 8 steps. When doing at step 9, we get 2722, which is larger than K . We stop and return $a = [0, 5, 1, 1, 2, 3, 1, 6, 2]$. It is $320/1789$.

Example 3. Let a be unknown rational number. We only know a bound of its denominator $N = 18$. According to theorem 7, Computing rational number a as follows: Take $K = N + 1$, yields $d = (2 * K + 2) * N * (N - 1) = 12240$. Assume that we use some numerical method to get an approximation 1.882434634 such that $|a - b| < 1/d$. Taking $\varepsilon = 1/K$, we recover number a by algorithm 1. We get $[1, 1, 7, 1, 1]$ by the first 5 steps. When doing at step

6, we get 41, which is larger than K . We stop and return $a = [1, 1, 7, 1, 1]$. It is $32/17$.

Example 4. This example is an application in obtaining exact factors from their approximations. Let $p = -16 - 56 * y - 48 * z + 64 * x^2 - 32 * x * y + 48 * x * z - 45 * y^2 - 96 * y * z - 27 * z^2$ be a polynomial. We want to use approximate method to get its exact factors over rational number field. First, we transform p to a monic polynomial as follows:

$$p = x^2 - \frac{1}{2}xy + \frac{3}{4}xz - \frac{45}{64}y^2 - \frac{3}{2}yz - \frac{27}{64}z^2 - \frac{7}{8}y - \frac{3}{4}z - \frac{1}{4}$$

the least common multiple of denominators of coefficients of polynomial $p(x, y, z)$ is 64, which is an upper bound[20] of denominators of coefficients of the monic factors of polynomial p . taking $K = 66$ yields $d = (2 * K + 2) * 65 * 64 = 557440$ and $\varepsilon = 1/K$. We use numerical methods to get its approximate factors as follows[20]:

$$\bar{g}_1 = 1.000000000000x + .6250000000067y + 1.124999999530z + .50000000000000$$

$$\bar{g}_2 = 1.000000000000x - 1.125000000015y - .3749999995480z - .50000000000000$$

the error of coefficients of $\bar{g}_1$ and $\bar{g}_2$ is less than $1/d$ by the numerical methods. According to theorem 7, taking ε in algorithm 1, we obtain two exact factors:

$$g_1 = x + \frac{5}{8}y + \frac{9}{8}z + \frac{1}{2}$$

$$g_2 = x - \frac{9}{8}y - \frac{3}{8}z - \frac{1}{2}$$

5 Conclusion

This paper builds a bridge spanning the gap between approximate computation and exact results. The exact results can be obtained by our algorithm as long as we get a bound N of absolute values of their denominators and their approximations with a error less than $1/((2N + 2)N(N - 1))$. Basing on our algorithm, we have succeed in obtaining exact factors of polynomials from their approximate factors. Our method can be applied in many aspect, such as proving inequality statements and equality statements, and computing resultants, etc. Thus we can take fully advantage of approximate methods to solve larger scale symbolic computation problems.

References

- [1] Beckermann, B., and Labahn, G., When are two polynomials relatively prime? *Journal of Symbolic Computation* 26 (1998), pp 677-689.
- [2] Corless R. M., Gianni P.M., Trager B.M. and Watt, S.M. The singular value decomposition for polynomial systems. In *International Symposium on Symbolic and Algebraic Computation* (Montreal, Canada, 1995), A Levelt, Ed., ACM pp. 195-207.
- [3] Corless, R.M., Giesbrecht, M.W., et al, Numerical implicitization of parametric hypersurfaces with linear algebra. In *proceeding of AISC2000, LNAI 1930*, pp.174-183.
- [4] Robert M. Corless, Mark W. Giesbrecht, et al, Towards factoring bivariate approximate polynomials, In *Proc. ISSAC 2001*, ACM press, pp.85-92
- [5] Robert M. Corless, Stephen M. Watt, and Lihong Zhi, QR Factoring to Compute the GCD of Univariate Approximate Polynomials *IEEE Transactions on Signal Processing*, 52(12) pp.3394-3402, 2004.
- [6] Robert M. Corless, Mark W. Giesbrecht, et al, Approximate polynomial decomposition. In *proceeding of ISSAC 1999*, S.S. Dooley, Ed., ACM pp 213-220.
- [7] Yong Feng, Yaohui Li, Checking RSC Criteria for Extended Dixon Resultant by Interpolation Method, *Proceedings of 7-th International Symposium on Symbolic and Numeric Algorithms for Scientific Computing*, Timisoara, Romania, September 25 - 29, 2005, IEEE Computer Society press pp48-51
- [8] Continued fraction, In: <http://www.answer.com>.
- [9] Galligo A., and Watt S. M. A numerical absolute primality test for bivariate polynomials In *proceeding of ISSAC 1997*. W. Küchlin, Ed. ACM, pp 217-224
- [10] Huang, Y., Wu, W., Stetter, H., and Zhi, L. Pseudofactors of multivariate polynomials. In *Proc. ISSAC'00(2000)*, ACM Press, pp.161-168.
- [11] Karmarkar N., and Lakshman Y. N., Approximate polynomial greatest common divisors and nearest singular polynomials. In *Proceeding of ISSAC 1996*, ACM, pp.35 -42.

- [12] Greg Reid , and Lihong Zhi, Solving Nonlinear Polynomial System via Symbolic-Numeric Elimination Method, In Proceedings of international conference on polynomial system solving. pp. 50-53. 2004
- [13] Zhi LIHONG and WU WENDA, Nearest Singular Polynomials Journal of Symbolic Computation, 26(6). pp.667-676, December 1998
- [14] Mou-Yan, Z., and Unbehausen, R. Approximate factorization of multi-variable polynomials. Signal Proces. 14(1988), 141-152.
- [15] Sasaki, T., Suzuki, M., et al., Approximate factorization of multivariate polynomials and absolute irreducibility testing. Japan J. Indust. Appl. Math. 8 (1991),357-375.
- [16] Sasaki, T., Saito T., and Hilano, T., Analysis of approximate factorization algorithm. Japan J. Indust. Appl. Math.9 (1992),351-368.
- [17] Tateaki Sasaki, Approximate multivariate polynomial factorization based on zero-sum relations. In Proc. ISSAC'2001, ACM Press, pp.284-291.
- [18] L. Yang, Jingzhong Zhang, and Xiaorong Hou, A Criterion of Dependency between Algebraic equation and its Application. Proceeding of IWMN'92, International Academic Publishers, pp 110-134, 1992.
- [19] Jingzhong Zhang, Yong Feng and Xijing Tang: Multivariate Polynomial factorization by Interpolation methods (extended abstract), Proc. the 7th Asian Symposium on Computer Mathematics (ASCM 2005), (Sung-il Pae, H. Park, eds.), Seoul, Dec.8-10, 2005.
- [20] Jingzhong Zhang, Yong Feng and Xijing Tang: Multivariate Polynomial factorization by Interpolation methods, submitted to China Sciences.